




Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)


Plan provided by  
Consonus Pharmacy

# Road to Recovery: Update



Information current as of 3.26.2021

## State Update: Ready to Activate!



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy






## Current State

### What we Know.....

- Keep going!
- Drop in case rate
- Reopening plan
- 5,000 Residents / 8,000 Staff left to vaccinate
- Consonus has 500 new recipients/week
- Pfizer is still available
- J&J – Nevada
- Moderna – still pending



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



## Current Consonus Stats Phase 2 (Beta):

Date of Last  
Clinic

3.19.2021

Date of Beta

3.1.2021



Agreements  
Received

85%

Facilities  
Activated

31

Phase 2  
Doses

559




CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy





# Policy Considerations: Community Based Care





## Step by Step Activation:



Activate:



## Activate

- Agreement
- Identify FC
- Identify Vaccinator
- FC Website Training
- Vaccinator Training
- Life Enrichment List
- Standing Orders
- Physician Order



Order



Receive



Vaccinate



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy





Thank you for your interest in participating in the COVID-19 Vaccine Continuity Program through Consonus Pharmacy. To reach activation status, please complete the following steps.

#### Step 1: Vaccination Agreement

- Return the signed vaccination agreement that was provided to you through DocuSign. If you participated in phase 1 clinics with Consonus, your agreement is already in place. If you are new to Consonus' vaccine program and did not receive your Vaccination Agreement, please contact your nurse or pharmacist consultant.

#### Step 2: Identify Facility Coordinator

- The Administrator, DNS /HWD or other leadership personnel within the facility should take on the Facility Coordinator (FC) role. The FC will be accountable and attest that requirements have been achieved for the Continuity Program at the facility. This person is responsible for ordering and securing the VaccPack as well as keeping the patient roster up to date in the Vaccine Administration Software (VAS). The FC will also manage an on call list to minimize waste of the vaccine at the facility site. Note, on call recipients must be within your local jurisdiction's current guidelines for COVID-19 vaccine eligibility.

#### Step 3: Identify Facility Vaccinator

- Secure healthcare provider(s) to administer vaccinations. Reconstitution may also be required, if utilizing the Pfizer product. Many states have expanded who can administer the COVID-19 vaccine during this national emergency. Please check your current state requirements on who can vaccinate.

#### Step 4: Facility Standing Orders

- Some states have standing orders for each vaccine product that allow healthcare professionals to administer the vaccine without individual patient orders. If your state does not have a standing order, the CDC has a template that may be utilized.

#### Step 5: Vaccine Allergic Reaction Box (VARB) Physician Order

- Each facility must purchase a VARB from the pharmacy to have onsite during vaccinations. Per the CDC guidance to administer the COVID vaccine, you must have 3 epinephrine auto injectors and an antihistamine (i.e. diphenhydramine). A link to a sample order is attached. Please have a prescriber sign and send back to pharmacy.

#### Step 6: Facility Coordinator and Vaccine Administrator Training

- Have VAS users view the following videos. Note, there have been several updates to the VAS since phase 1. For this reason, regardless if you have participated in phase 1 clinics, you are required to watch the appropriate trainings for your role.

##### Facility Coordinator:

- ["Facility Coordinator VAS Training Video"](#)

##### Vaccinator:

- ["Vaccine Administrator VAS Training"](#)
- ["Vaccine Prep and Administration"](#) (product specific)

#### Step 7: Activation Confirmation or Training Access

- Once you have completed all steps, please fill out the ["Ready to Activate"](#) form to validate all steps have been completed. A vaccine coordinator will then contact you with your first order date.



## Activation Check List:

### Ready Set GO! Activation Check List

<input type="checkbox"/>	Vaccination Agreement	
<input type="checkbox"/>	Identify Facility Coordinator	
<input type="checkbox"/>	Identify Facility Vaccinator	
<input type="checkbox"/>	Facility Standing Order	Check state requirements <a href="#">Washington State Standing Orders</a> <a href="#">CDC Standing Order Pfizer</a> <a href="#">CDC Standing Order Moderna</a>
<input type="checkbox"/>	Vaccine Allergic Reaction Box Order	<a href="#">WA Pharmacy Customer</a> <a href="#">OR Pharmacy Customer</a> <a href="#">CA Pharmacy Customer</a> <a href="#">NV Pharmacy Customer</a>
<input type="checkbox"/>	Facility Coordinator Training	<a href="#">Facility Coordinator VAS Training Video</a> <a href="#">Review Training Guide</a>
<input type="checkbox"/>	Vaccinator Training	<a href="#">Vaccine Administrator VAS Training</a> <a href="#">Vaccine Prep and Administration</a> (product specific) <a href="#">Review Training Guide</a>
<input type="checkbox"/>	Fill Out "Ready to Activate" Form	<a href="#">Ready to Activate</a>

Please note, Consonus vaccine supply may vary based on state and federal allocation. Fluctuating vaccine supply may delay facilities preferred vaccination schedule.



## Step 1: Vaccination Agreement



- Phase 1 participants – Agreement already in place.
- Parent company agreements may be in place.
- If you are unsure if you have an agreement or need one, please reach out to you consultant pharmacist or nurse.

DocuSign®



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY  
*Ready, Set, Go*






## Step 2: Identify Facility Coordinator

### Who:

- Leadership

### Responsibilities:


- Attest to requirements
- Maintain roster
- Order vaccine
- Coordinate delivery
- Maintain security
- Report end of clinic inventory
- Maintain life enrichment list







## Step 3: Identify Facility Vaccinator



### Responsibilities:

- Clinical screening
- Vaccination
- Documentation in VAS
- CDC vaccination card
- Post vaccine monitoring



## Step 4: Clinical Standing Orders for Administration

### Moderna COVID-19 Vaccine

Standing Orders for Administering Vaccine  
to Persons 18 Years of Age and Older



**Note:** For more information/guidance, please contact the immunization program at your state or local health department or the appropriate state body (e.g., state board of medical/nursing/pharmacy practice).

#### Purpose

- To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

#### Policy

- Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

#### Procedure

Assess persons 18 years of age and older for vaccination with Moderna COVID-19 Vaccine based on the following criteria:

- No complete 2-dose COVID-19 vaccination history, regardless of brand. If 2 doses of a same-brand or mixed-brand series have been administered, no additional doses are recommended.
- If the recipient has received 1 previous dose of Moderna COVID-19 Vaccine, a second dose of the same brand should be administered.
- If the vaccine product previously given cannot be determined or is no longer available, any mRNA COVID-19 vaccine product may be administered at least 28 days after the first dose.
- This vaccine is administered in a 2-dose series. Separate doses by at least 28 days, preferably within 6 weeks.\*

- Moderna COVID-19 Vaccine should not be administered at the same time as other vaccines. Separate Moderna COVID-19 Vaccine from other vaccines by 14 days before or after the administration of Moderna COVID-19 Vaccine.<sup>†</sup>

- Moderna COVID-19 Vaccine should be deferred for at least 90 days for persons who received passive antibody therapy (monoclonal antibodies or convalescent plasma) as part of COVID-19 treatment.

- If the recipient has a history of dermal filler use, advise them to contact their healthcare provider for evaluation if they develop swelling at or near the dermal filler site following vaccination.
- Screen for contraindications and precautions.

#### Contraindications:

- Severe allergic reaction (e.g., anaphylaxis) to a previous dose or component of either mRNA COVID-19 vaccine.
- Immediate allergic reaction<sup>‡</sup> of any severity to a previous dose or component of an mRNA COVID-19 vaccine (including polyethylene glycol [PEG]). See Table 1 of vaccine components on page 3.
- Immediate allergic reaction of any severity to polysorbate (due to potential cross-reactive hypersensitivity with the vaccine ingredient PEG).

#### Precautions:

- History of an immediate allergic reaction to any other vaccine or therapies [excluding subcutaneous immunotherapy of allergies, i.e., "allergy shots"] not related to a component of mRNA COVID-19 vaccines or polysorbate).
- Moderate to severe acute illness.

- Provide all recipients with a copy of the current federal Emergency Use Authorization (EUA) Fact Sheet for Recipients and Caregivers.
- Prepare to administer the vaccine.

Sex and Weight of Patient	Needle Gauge	Needle Length	Injection Site <sup>§</sup>
Female or male fewer than 130 lbs	22–25	½"–1"	Deltoid muscle of arm
Female or male 130–152 lbs	22–25	1"	Deltoid muscle of arm
Female 152–200 lbs	22–25	1–1½"	Deltoid muscle of arm
Male 153–260 lbs	22–25	1–1½"	Deltoid muscle of arm
Female 200+ lbs	22–25	1½"	Deltoid muscle of arm
Male 260+ lbs	22–25	1½"	Deltoid muscle of arm

\*Administer the second dose as close to the recommended interval (28 days) as possible. If the second dose is not administered within 6 weeks of the first dose, the series does not need to be restarted. Doses inadvertently administered less than 28 days apart do not need to be repeated.

†However, mRNA COVID-19 and other vaccines may be administered within a shorter period in situations where the benefits of vaccination are deemed to outweigh the potential unknown risks of vaccine coadministration (e.g., tetanus toxoid-containing vaccination as part of wound management, rabies vaccination for post-exposure prophylaxis, measles or hepatitis A vaccination during an outbreak) or to avoid barriers or delays to mRNA COVID-19 vaccination.

‡For the purpose of this guidance, an immediate allergic reaction is defined as any hypersensitivity-related signs or symptoms such as urticaria, angioedema, respiratory distress (e.g., wheezing, stridor), or anaphylaxis that occur within 4 hours following exposure to a vaccine or medication.

§Alternatively, the anterolateral thigh also can be used.

§Some experts recommend a 5/8-inch needle for men and women who weigh less than 130 pounds. If used, skin must be stretched tightly (do not bunch subcutaneous tissue).

02/11/2021 CS321571.H

1

### Pfizer-BioNTech COVID-19 Vaccine

Standing Orders for Administering Vaccine  
to Persons 16 Years of Age and Older



**Note:** For more information/guidance, please contact the immunization program at your state or local health department or the appropriate state body (e.g., state board of medical/nursing/pharmacy practice).

#### Purpose

- To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

#### Policy

- Where authorized under state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess and vaccinate persons who meet the criteria in the "Procedure" section below without the need for clinician examination or direct order from the attending provider at the time of the interaction.

#### Procedure

- Assess persons 16 years of age and older for vaccination with Pfizer-BioNTech COVID-19 Vaccine based on the following criteria:

- No complete 2-dose COVID-19 vaccination history, regardless of brand. If 2 doses of a same-brand or mixed-brand series have been administered, no additional doses are recommended.
- If the recipient has received 1 previous dose of Pfizer-BioNTech COVID-19 Vaccine, the second dose of the same brand should be administered.
- If the vaccine product previously given cannot be determined or is no longer available, any mRNA COVID-19 vaccine product may be administered at least 28 days after the first dose.
- This vaccine is administered in a 2-dose series. Separate doses by at least 28 days, preferably within 6 weeks.\*

- Pfizer-BioNTech COVID-19 Vaccine should not be administered at the same time as other vaccines. Separate Pfizer-BioNTech COVID-19 Vaccine from other vaccines by 14 days before or after the administration of Pfizer-BioNTech COVID-19 Vaccine.<sup>†</sup>

- Pfizer-BioNTech COVID-19 Vaccine should be deferred for at least 90 days for persons who received passive antibody therapy (monoclonal antibodies or convalescent plasma) as part of COVID-19 treatment.

- Screen for contraindications and precautions.

#### Contraindications:

- Severe allergic reaction (e.g., anaphylaxis) to a previous dose or component of either mRNA COVID-19 vaccine.
- Immediate allergic reaction<sup>‡</sup> of any severity to a previous dose or component of an mRNA COVID-19 vaccine (including polyethylene glycol [PEG]). See Table 1 of vaccine components on page 3.
- Immediate allergic reaction of any severity to polysorbate (due to potential cross-reactive hypersensitivity with the vaccine ingredient PEG).

#### Precautions:

- History of an immediate allergic reaction to any other vaccine or therapies [excluding subcutaneous immunotherapy of allergies, i.e., "allergy shots"] not related to a component of mRNA COVID-19 vaccines or polysorbate).
- Moderate to severe acute illness.

- Provide all recipients with a copy of the current federal Emergency Use Authorization (EUA) Fact Sheet for Recipients and Caregivers.

- Prepare to administer the vaccine.

Sex and Weight of Patient	Needle Gauge	Needle Length	Injection Site <sup>§</sup>
Female or male fewer than 130 lbs	22–25	½"–1"	Deltoid muscle of arm
Female or male 130–152 lbs	22–25	1"	Deltoid muscle of arm
Female 152–200 lbs	22–25	1–1½"	Deltoid muscle of arm
Male 153–260 lbs	22–25	1–1½"	Deltoid muscle of arm
Female 200+ lbs	22–25	1½"	Deltoid muscle of arm
Male 260+ lbs	22–25	1½"	Deltoid muscle of arm

\*Administer the second dose as close to the recommended interval (21 days) as possible. If the second dose is not administered within 6 weeks of the first dose, the series does not need to be restarted. Second doses inadvertently administered less than 21 days apart do not need to be repeated.

†However, mRNA COVID-19 and other vaccines may be administered within a shorter period in situations where the benefits of vaccination are deemed to outweigh the potential unknown risks of vaccine coadministration (e.g., tetanus toxoid-containing vaccination as part of wound management, rabies vaccination for post-exposure prophylaxis, measles or hepatitis A vaccination during an outbreak) or to avoid barriers or delays to mRNA COVID-19 vaccination.

‡For the purpose of this guidance, an immediate allergic reaction is defined as any hypersensitivity-related signs or symptoms such as urticaria, angioedema, respiratory distress (e.g., wheezing, stridor), or anaphylaxis that occur within 4 hours following exposure to a vaccine or medication.

§Alternatively, the anterolateral thigh also can be used.

§Some experts recommend a 5/8-inch needle for men and women who weigh less than 130 pounds. If used, skin must be stretched tightly (do not bunch subcutaneous tissue).

02/11/2021 CS321570.H

1



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY  
Ready, Set, Go





## Step 5: Requirement to Give Vaccine



- Epinephrine in prefilled autoinjector
- Antihistamine (Benadryl)
- Access to call 911



- First Aid Kit



- Stethoscope



- Blood pressure cuff



- Timing device for pulse



- Oxygen



- Pulse Oximeter







# Step 5: Vaccine Allergic Reaction Box (VARB)



- Epinephrine in prefilled autoinjector (3)
- Antihistamine (diphenhydramine)







## Step 6: Training



### Required Training:

- Facility Coordinator Vaccine Administration Software Video
- Knowledge of current jurisdiction for vaccinating



### Required Training:

- “Vaccine Administrator VAS Training” Video
- “Vaccine Prep and Administration” Video
- Knowledge of current jurisdiction for vaccinating







# Step 7: Ready to Activate?

- Fill out “Ready to Activate” form
- After completion of activation steps
- Attest that all steps have been achieved
- Consonus to verify
- Confirmation email sent to facility coordinator

### Ready to Activate Form

Please complete and submit the following form to indicate readiness for activation with the Consonus Vaccine Continuity Program.

After submission of this form, Consonus will contact the Facility Coordinator named below to coordinate activation and notify you of the earliest date you can place a vaccine order.

Please note, Consonus vaccine supply may vary based on state and federal allocation. Fluctuating vaccine supply may delay facilities' preferred vaccination schedule.

\* Required

1. Facility Name \*

Enter your answer

2. Facility Coordinator Name \*

Enter your answer

3. Facility Coordinator Email Address \*

Enter your answer

5. All of the following items have been completed: \*

☐ Signed Vaccine Continuity Program agreement sent to Consonus

☐ Signed prescription for epinephrine under the name of the facility sent to Consonus

☐ Standing order to administer vaccine obtained

6. The Facility Coordinator has reviewed all of the following training materials related to the vaccine ordering process and Vaccine Administration Software (VAS): \*

Training materials can be accessed here:  
<https://confluence.marquiscompanies.com/pages/viewpage.action?pageId=54887559>

☐ Watched Facility Coordinator VAS training video

☐ Watched Vaccine Administrator VAS training video

☐ Reviewed VAS guide

☐ Reviewed Consonus Pharmacy Vaccine Delivery Schedule

Submit

Never give out your password. Report abuse




## Moving from Clinics to Continuity:




Activate

- Agreement
- Identify FC
- Identify Vaccinator
- FC Website Training
- Vaccinator Training
- Physician Order
- Standing Orders
- Life Enrichment List



Order

- Maintain patient's roster
- e-Consent
- EUA
- 2 Days in Advance
- Rx Verification



Receive



Vaccinate



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



## Ordering Key Points:




Activation completed  
and confirmed




Roster uploaded prior to  
ordering



Order 2 days in Advance



Vaccinator Scheduled




Shipped Monday – Friday Only



- **Your clinics** should be  
scheduled **Tuesday - Saturday**



Know State Phase Criteria

# Ordering the VaccPack:





My Patients

Upcoming Clinics

Vaccine Inventory

Manage Accounts

My Clinics

Help

Order Pfizer

Inventory Report Out

Patient Corrections

My Patients at

Total Number of Residents

22

Obtained (82%)

Refused (0%)

Needed (18%)

Stop - Do Not Give (0%)

Total Number of Staff

3

Obtained (67%)

Refused (0%)

Needed (33%)

Stop - Do Not Give (0%)

Vaccination Progress

27

1st Dose (11%)

None (44%)

Booster (44%)

Hope Village Patients

Search...

FILTER


EXPORT PATIENTS

ADD PATIENT

!	First Name	Last Name	Patient Type	First Dose	Booster Dose	Scheduled	Consent Status
	Another	Patient3	Resident	12/20/2020	12/23/2020	Yes	Obtained
	Cool	Guy	Resident	02/02/2021	02/26/2021	Yes	Obtained

Consonus Healthcare  
consonushealth.com


Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY  
Ready, Set, Go



# Ordering the VaccPack:



## Pfizer BioNTech VACCINE ORDER FORM

The survey will take approximately 4 minutes to complete.

\* Required

1. Servicing Pharmacy \*

- ☐ Consonus Pharmacy
- ☐ Community Pharmacy

2. Scheduled Date for Vaccinations: \*

Please input date in format of M/d/yyyy



3. I attest that Vaccine Administrators and I have read, viewed, and understand the training documents and videos shown under the "Training Materials" tab on the Consonus/Community Vaccine Administration Site (CVAS) which discuss the storage and handling, preparation and dilution, and administration of the Pfizer-BioNTech COVID-19 vaccine: \*

Link to Training Materials:

<https://confluence.marquiscompanies.com/display/KB/Vaccine+Prep+and+Administration+Library>

- ☐ I attest Vaccine Administrators and I have reviewed and understand the above training materials



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



# Ordering the VaccPack:

## Roster Verified by Consonus





## Moving from Clinics to Continuity:




### Activate

- Agreement
- Identify FC
- Identify Vaccinator
- FC Website Training
- Vaccinator Training
- Physician Order
- Standing Orders
- Life Enrichment List



### Order

- Maintain patient's roster
- e-Consent
- EUA
- 2 Days in Advance
- Rx Verification



### Receive

- Consistent Location
- Secure
- Do Not Open



### Vaccinate




CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



# Introducing the VaccPack:




- Vaccine
- Diluent – 0.9% NS Vial
- Vaccine Dilution Instructions
- Syringe(s)
- Alcohol Swabs
- CDC Cards
- Band-Aides

Product is to stay in pharmacy packaging until ready to vaccinate.



# Receiving:



## Routine Delivery:

- Use within 24 hours of receipt
- Product is to stay in pharmacy packaging until ready to vaccinate.

## Moving from Clinics to Continuity:



### Activate

- Agreement
- Identify FC
- Identify Vaccinator
- FC Website Training
- Vaccinator Training
- Physician Order
- Standing Orders
- Life Enrichment List



### Order

- Maintain patient's roster
- e-Consent
- EUA
- 2 Days in Advance
- Rx Verification



### Receive

- Central Location
- Secure
- Do Not Open



### Vaccinate

- Allergic Reaction Box
- Vaccination Kit
- Screen
- Monitor
- Inventory Report



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY

Ready, Set, Go



# Screening Protocols Remain in Place

## Pre-Vaccination Form for Pfizer-BioNTech COVID-19 Vaccine



Information for Healthcare Professionals about the Pre-Vaccination Form for Pfizer-BioNTech COVID-19 Vaccine.

For additional information on COVID-19 vaccine recommendations, see:  
<https://www.cdc.gov/vaccines/covid-19/info-by-product/pfizer/covid-19-considerations.html>  
For additional information on ACIP general recommendations, see:  
<https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html>

### Are you feeling sick today?

There is no evidence that acute illness reduces vaccine efficacy or increases vaccine adverse events. However, as a precaution with moderate or severe acute illness, all vaccines should be delayed until the illness has improved. Mild illnesses (e.g., upper respiratory infections, diarrhea) are NOT contraindications to vaccination. Do not withhold vaccination if a person is taking antibiotics. Vaccination of persons with current SARS-CoV-2 infection should be deferred until the person has recovered from acute illness and they can discontinue isolation. While there is no minimum interval between infection and vaccination, current evidence suggests reinfection is uncommon in the 90 days after initial infection. Persons with documented acute SARS-CoV-2 infection in the preceding 90 days may delay vaccination until near the end of this period, if desired.

### Have you ever received a dose of COVID-19 vaccine?

Two doses of the same COVID-19 vaccine product are recommended. Check medical records, immunization information systems, and vaccination record cards to help determine the initial product received. Those who received a trial vaccine should consult with the trial sponsors to determine if it is feasible to receive additional doses.

Have you ever had a severe allergic reaction (e.g., anaphylaxis) to something? For example, a reaction for which you were treated with epinephrine or EpiPen®, or for which you had to go to the hospital? Allergic reactions, including severe allergic reactions, NOT related to vaccines or injectable therapies (e.g., food, pet, venom, environmental, or latex allergies; oral medications) are NOT a contraindication or precaution to vaccination with currently authorized COVID-19 vaccine. HOWEVER, individuals who have had severe allergic reactions to something, regardless of cause, should be observed for 30 minutes after vaccination. All other persons should be observed for 15 minutes.

### Was the severe allergic reaction after receiving a COVID-19 vaccine?

History of severe allergic reaction (e.g., anaphylaxis) to a previous dose or component of the COVID-19 vaccine product being offered is a contraindication to that COVID-19 vaccine.

### Was the severe allergic reaction after receiving another vaccine or another injectable medication?

A history of mild allergic reaction to a vaccine or injectable therapy is not a precaution to vaccination. History of severe allergic reaction (e.g., anaphylaxis) to another vaccine or a component of another vaccine OR anaphylactic reaction to any other injectable medication is a precaution to currently authorized COVID-19 vaccine. Vaccine may be given, but counsel patients about unknown risks of developing a severe allergic reaction and balance these risks against the benefits of vaccination. These individuals should be observed for 30 minutes after vaccination.

### Do you have a bleeding disorder or are you taking a blood thinner?

COVID-19 vaccine may be given to these patients, if a physician familiar with the patient's bleeding risk determines that the vaccine can be administered intramuscularly with reasonable safety. ACIP recommends the following technique for intramuscular vaccination in patients with bleeding disorders or taking blood thinners: a fine-gauge needle (23-gauge or smaller caliber) should be used for the vaccination, followed by firm pressure on the site, without rubbing, for at least 2 minutes.

### Have you received passive antibody therapy as treatment for COVID-19?

Based on the estimated half-life of monoclonal antibodies or convalescent plasma as part of COVID-19 treatment, as well as evidence suggesting that reinfection is uncommon in the 90 days after initial infection, vaccination should be deferred for at least 90 days, as a precautionary measure until additional information becomes available, to avoid interference of the antibody treatment with vaccine-induced immune responses.

### » Considerations

Immunocompromise is not a contraindication to current COVID-19 vaccine, including those with cancer, leukemia, HIV/AIDS and other immune system problems or taking medication that affects their immune systems. However, patients should be informed that the vaccine might be less effective than in someone who is immunocompetent.

Pregnancy is not a contraindication to current COVID-19 vaccine. While there are currently no available data on the safety of COVID-19 vaccines in pregnant people, studies and results are expected soon. Pregnant people may choose to get vaccinated. Observational data demonstrate that while the absolute risk is low, pregnant people with COVID-19 have an increased risk of severe illness.

Lactation is not a contraindication to current COVID-19 vaccine. Lactating people may choose to be vaccinated. There is no data available for lactating people on the effects of mRNA vaccines.

12/16/20 C031620-6

2

## Pre-Vaccination Form for Pfizer-BioNTech COVID-19 Vaccine



### For vaccine recipients:

The following questions will help us determine if there is any reason you should not get the COVID-19 vaccine today. If you answer "yes" to any question, it does not necessarily mean you should not be vaccinated. It just means additional questions may be asked. If a question is not clear, please ask your healthcare provider to explain it.

Patient Name \_\_\_\_\_

Age \_\_\_\_\_

	Yes	No	Don't know
1. Are you feeling sick today?			
2. Have you ever received a dose of COVID-19 vaccine? If yes, which vaccine product? <input type="checkbox"/> Pfizer <input type="checkbox"/> Another product _____			
3. Have you ever had a severe allergic reaction (e.g., anaphylaxis) to something? For example, a reaction for which you were treated with epinephrine or EpiPen®, or for which you had to go to the hospital?			
• Was the severe allergic reaction after receiving a COVID-19 vaccine?			
• Was the severe allergic reaction after receiving another vaccine or another injectable medication?			
4. Do you have a bleeding disorder or are you taking a blood thinner?			
5. Have you received passive antibody therapy as treatment for COVID-19?			

Form completed by \_\_\_\_\_ Date \_\_\_\_\_

Form reviewed by \_\_\_\_\_ Date \_\_\_\_\_

Adapted with appreciation from the Immunization Action Coalition (IAC) screening checklist

12/16/20 C031620-6

1



CONSONUS  
PHARMACY



Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY  
Ready, Set, Go

# Report Inventory:



My Patients

Upcoming Clinics

Vaccine Inventory

Manage Accounts

My Clinics

Help

Order Pfizer

Inventory Report Out

Patient Corrections

My Patients at

Total Number of Residents

22

Obtained (82%)

Refused (0%)

Needed (18%)

Stop - Do Not Give (0%)

Total Number of Staff

3

Obtained (67%)

Refused (0%)

Needed (33%)

Stop - Do Not Give (0%)

Vaccination Progress

27

1st Dose (11%)

None (44%)

Booster (44%)

Hope Village Patients


Search...

FILTER

EXPORT PATIENTS


ADD PATIENT

!	First Name	Last Name	Patient Type	First Dose	Booster Dose	Scheduled	Consent Status
	Another	Patient3	Resident	12/20/2020	12/23/2020	Yes	Obtained
	Cool	Guy	Resident	02/02/2021	02/26/2021	Yes	Obtained



Consonus Healthcare  
consonushealth.com


Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY  
Ready, Set, Go



# Report Inventory:



## Post-Clinic Inventory Report Out

Facility Coordinator to complete the below Inventory Report

\* Required

1. Date of Vaccine Adminsitration \*

Please input date in format of M/d/yyyy



2. Name of Facility (Include ALF, SNF, MC in Name) \*

Enter your answer

3. Vaccine Manufacturer: \*

☐ Pfizer-BioNTech

☐ Moderna

4. Number of Vials Diluted: \*

# COVID Vaccines Authorized for Use:



- Pfizer
- Moderna
- Janssen

The best vaccine is  
the one readily  
available to you!

1

Goal

2








Efficacy

3

Safety









## Storage and Handling:

				
	Age	18 yo	16 yo	18 yo
	Between doses	28 days	21 Days	Single Dose
	Room Temp (unopened vials)	12 hours	2 hours	12 hours
	In use vials	6 hours after puncture	6 hours after dilution	2 hours- rm temp 6 hours - refrigerator




# Product Comparison:

				
	Dilution	None, ready to use	Dilute with saline	None, ready to use
	Doses/Vial	10	6	5
	Dose	0.5 ml IM	0.3ml IM	0.5 ml IM



#GetVaccinated:



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY  
*Ready, Set, Go*

# Pharmacy Updates:

Pharmacy  
lowering infection  
control policy to  
level 2!

April 1, 2021



---

Going back to pre-COVID delivery methods



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy





## Pharmacy Updates:



Continued Consulting  
Service....  
but smiles in person!



CONSONUS  
PHARMACY

Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy



ROAD TO  
RECOVERY  
*Ready, Set, Go*

## Best Case Schedule:



### Get Ready!

- ✓ Week 1 – 2 Pilot facilities
- ✓ Week 2 – 21 facilities
- ~~Week 3 – Continue SNF + 4 ALF's~~
- ✓ Week 4 <sup>3</sup> – All SNF Customers + Additional <sup>ALL</sup> ALF's



# The Goal:



Maintain  $\geq 75\%$  full vaccination protection within LTC facilities.



Decrease community positivity rates.




Reopen! Bring community and families back together.



Maximize product, minimizing waste

## Get Started?



1

Complete Activation Checklist

2

Order VaccPack!



CONSONUS  
PHARMACY


Consonus Healthcare  
[consonushealth.com](https://consonushealth.com)

Plan provided by  
Consonus Pharmacy





# Questions?



- Check our website for the latest information
- This presentation will be sent out and available on the website **Monday**
- FAQ Friday answering your commonly asked questions
- When we have information, you will hear from us